



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

ture of their antennæ, maxillæ, labium, etc. Of those two groups we possess, as far as I am aware, only two good descriptions of larvæ: Mulsant's of the larva of *Geotrupes stercorarius* and Chapuis' of the larva of *Trox carolinus*. If we were to base our opinion upon these two descriptions only, it would incline in favor of a relationship of our larva with the *Trogidæ*, rather than with the *Geotrupidæ*. Chapuis' description of the larva of *Trox carolinus* agrees quite well with our larva; the description of the labium especially (lèvre inférieure formée d'un menton et d'une pièce palpigère fondus en un real corps allongé) seems to indicate a structure somewhat analogous to that in our larva. On the contrary, Mulsant's description of the larva of *Geotrupes* disagrees with ours in several points. The antennæ are said to be four-jointed, the maxillæ to have two almost cylindrical lobes, the legs are described as bilobed at the end, etc. Finally, if it be true that the larva of *Geotrupes* has only two pairs of well developed legs, the third being almost atrophied, as Frisch (but not Mulsant) describes it, this would constitute another important difference.

Altogether, the materials at hand for comparison are too meagre for a final decision upon the relationship of the larva.

Before concluding I would advert once more to the singular fact that the shape of the mandibles changes after moulting. A similar peculiarity has been already observed among larvæ of other orders of insects. That the shape of the earlier mandibles is merely due to its being more worn is a supposition which will hardly be entertained by any one who has compared the two mandibles.


EXPLANATION OF THE FIGURES.

Fig. 1.—Larva from the side, natural size.

Fig. 2.—Head from above.

Fig. 3.—Labium, maxilla, and in front of them, the mandible *after moulting*.

Fig. 4.—Leg.



On the CUPESIDÆ of North America.

BY JOHN L. LECONTE, M. D.

PRIACMA, n. g.

Since I have seen the Australian genus *Omma*, I recognize in *Cupes serrata* (Lec. Proc. Acad. Nat. Sc. Phila., 1861, 351) a distinct genus, intermediate between that and *Cupes*. From the latter it differs by the antennæ less approximate at base, shorter and stouter,

scarcely half as long as the body, with the joints triangular and narrower at the base, the 1st as long as but stouter than the 3d. The eyes are much smaller, though nearly smooth; the under surface of the head is quite different; gular sutures distant, curved slightly outwards, gular angles not porrected, but broad and truncate; mentum more prominent, maxillæ more exposed, maxillary palpi longer, last joint elongate, cylindrical, truncate. These differences were partly indicated by me in the remarks appended to the description of the species. The color is mottled gray; head with four subacute tubercles, but not lobed, front concave. Prothorax nearly square, with front angles prominent; elytra more convex, alternate interspaces feebly convex and a little uneven; side margins strongly toothed towards the tip.

Cupes lobiceps.—Body elongate, slender, pale gray, slightly mottled with darker; antennæ compressed, serrate for the lower half, two-thirds as long as the body; head deeply channelled and transversely impressed, thus divided into two large posterior tubercles or lobes, and two smaller frontal ones; there is also a lateral lobe behind the eye, separated by a deep groove, eyes large; prothorax suddenly narrowed in front, with the sides strongly elongated and a large median elevation which has a deep rhomboidal excavation; elytra with rows of moderate sized quadrate punctures, 3d interspace more convex, 5th, 6th and 7th forming an obtuse costa, 2d and 4th flat. Length .45 inch; 11 mm.

San Diego, California. A specimen kindly given to me by Mr. Ulke. This species resembles the figure of *C. Latreillei* Solier (Lacordaire, Gen. Col. Atlas, pl. 47, fig. 2) in the form of the antennæ. It differs conspicuously from our other species by the large posterior lobes of the head and the different sculpture of the prothorax. The following table will enable them to be easily recognized:—

- A.—Antennæ serrate; gular sutures converging behind; eyes large. Cinereous mottled; head with four tubercles on the upper surface and two lateral; prothorax with elevated margins and disc..... **1. lobiceps.**
- B.—Gular sutures parallel.
- a.—Antennæ as long as the body, filiform; eyes large. Cinereous mottled; head with four feebly separated tubercles; prothorax transverse, sides not elevated..... **2. concolor.**
- b.—Antennæ nearly cylindrical, one-half as long as the body; eyes small. Black; head with fulvous hair; with two frontal and four occipital tubercles, the middle ones smaller and less elevated..... **3. capitata.**